

Safety valves for photosynthesis

Krishna K Niyogi

Recent studies have provided new insights into the ways that plants may dissipate excess photons and electrons, thereby protecting the photosynthetic apparatus against light-induced damage. These 'safety valves' include nonphotochemical mechanisms for quenching excited chlorophylls, as well as alternative electron acceptors such as oxygen.

Addresses

Department of Plant and Microbial Biology, University of California, Berkeley, California 94720-3102, USA;
e-mail: niyogi@nature.berkeley.edu

Current Opinion in Plant Biology 2000, 3:455–460

1369-5266/00/\$ – see front matter

© 2000 Elsevier Science Ltd. All rights reserved.

Abbreviations

Chl	chlorophyll
im	<i>immutans</i>
LHC	light-harvesting complex
NPQ	nonphotochemical quenching
qE	Δ pH-dependent quenching
qI	photoinhibitory quenching
PS II	photosystem II
PsbS	PS II subunit S
Rubisco	ribulose-1,5-bisphosphate carboxylase/oxygenase

Introduction

The photosynthetic apparatus in plants performs the essential function of converting light energy into chemical energy that is used for CO₂ fixation and other assimilatory reactions, but this complex machinery is susceptible to light-induced damage caused by the inevitable generation of reactive intermediates and by-products. Diurnal and seasonal fluctuations in light intensity, as well as environmental stresses such as cold, drought, salinity, and nutrient deficiency, that limit CO₂ fixation can result in the absorption of more light energy than can be utilized productively by photosynthesis. Under these conditions, safe dissipation of excess photons and electrons is necessary to protect the photosynthetic apparatus from light-induced damage.

Maintenance of photosynthesis is critical for plant fitness and survival, as evidenced by the light sensitivity of many photosynthetic mutants and antisense plants, and therefore plants have evolved numerous photoprotective mechanisms [1*]. These include several antioxidant molecules and enzymes [1*], repair processes for lipid peroxidation [2] and damaged photosystem II (PS II) reaction centers [3], and even systemic signaling and acclimation processes [4]. This short review will focus on the various 'safety valves' that may be involved in photoprotection through dissipation of excess photons and electrons (Figure 1).

The constitutive triplet chlorophyll valve

Light is absorbed by chlorophyll (Chl) and carotenoid molecules that are bound to light-harvesting complex

(LHC)-proteins in the thylakoid membranes of chloroplasts [5], resulting in singlet state excitation of the pigment molecules. Excitation energy is then transferred to reaction centers to drive electron transport, which oxidizes H₂O, reduces NADP⁺, and generates a Δ pH that is harnessed for ATP synthesis. In some plants, leaf movements allow the plant to adjust the angle of a leaf relative to the sun in order to optimize (or minimize) light absorption. However, during the course of a typical day most plants are simply unable to regulate light absorption on a timescale that is fast enough to cope with the diurnal and other rapid fluctuations in light intensity that occur in natural environments. Instead of changing the capacity for light absorption, plants rely on mechanisms for dissipating photons that have already been absorbed.


Absorption of excess photons can potentially cause accumulation of excitation energy in the LHCs and thereby increase the lifetime of singlet Chl (¹Chl). Through a physical process known as intersystem crossing, ¹Chl can be converted into triplet Chl (³Chl), a long-lived state that is incapable of initiating photosynthetic electron transfer. Generation of ³Chl results in the dissipation of absorbed photons because ³Chl is efficiently quenched by energy transfer to carotenoids, which are closely associated with Chls in the photosynthetic apparatus. It has been estimated that between 4% and 25% of the photons absorbed by PS II can be dissipated via this triplet valve. The proportion of photons dissipated in this way is correlated to the average lifetime of ¹Chl [6]. However, because the lifetime of ¹Chl is kept relatively low and constant by regulated de-excitation (see below), dissipation via the triplet valve is usually limited to the lower end of the estimated 4–25% range.

Although ³Chl itself is not harmful, it can interact with ground state molecular O₂, which normally exists as a triplet, converting it into singlet oxygen (¹O₂), a highly reactive oxygen species. Production of ¹O₂ has been detected in leaves illuminated with intense light [7], and it can directly damage the proteins, pigments, and lipids that comprise the photosynthetic apparatus. Quenching of ¹O₂ is also accomplished by carotenoids, which explains why these pigments are essential for photosynthesis in the presence of oxygen.

Rapidly inducible dissipation of excess photons

Whereas the de-excitation of ¹Chl via the triplet valve is basically an unregulated, constitutive process, plants and most eukaryotic algae also possess a rapidly inducible nonphotochemical quenching (NPQ) mechanism, termed Δ pH-dependent quenching (qE), for the harmless thermal dissipation of excess absorbed photons in PS II [8] (Figure 1). The qE mechanism can be considered as a feedback-regulated safety valve for photosynthesis because it is controlled primarily by the magnitude of the thylakoid Δ pH that is generated by photosynthetic electron transport.

Figure 1



Schematic depiction of safety valves for excess photons and electrons that are discussed in the text. Thermal dissipation of excess photons is accomplished by the triplet chlorophyll valve (³Chl), inducible qE, and

sustained qI. Oxygen-dependent alternative electron sinks include photorespiratory metabolism, photoreduction of O₂ in the water–water cycle, and a chlororespiratory alternative oxidase.

Absorption of excess photons causes the buildup of a high ΔpH , and the resulting decrease in lumen pH is essential for qE [8]. The lowering of the lumen pH activates violaxanthin de-epoxidase, a luminal enzyme that converts violaxanthin to zeaxanthin as part of the xanthophyll cycle [9]. A strong correlation between zeaxanthin accumulation and qE has been demonstrated for a wide variety of plants under both natural and laboratory conditions [10]. Low lumen pH may also result in the protonation of one or more protein(s) in the LHC that is associated with PS II [8]. Together, binding of protons and zeaxanthin is postulated to cause a conformational change, which can be assayed as an absorbance change at 535 nm in leaves or isolated thylakoids, that effectively switches the LHC into a state in which efficient nonphotochemical de-excitation of ¹Chl and concomitant thermal dissipation of the excitation energy occur. The switch to thermal dissipation is accompanied by a discrete change in Chl-fluorescence-lifetime distributions [11]. The conformational change and the conversion of the LHC into a quenched state are rapidly reversible when light is no longer excessive and the magnitude of the ΔpH decreases.

The recent isolation and characterization of mutants are providing new insights into qE [12–15,16••]. For example, the role of xanthophylls in qE has been addressed using mutations that affect xanthophyll synthesis. The *npq1* mutant of the common laboratory weed, *Arabidopsis thaliana*, is defective in the gene encoding violaxanthin de-epoxidase, so it is unable to convert violaxanthin to zeaxanthin in the xanthophyll cycle [13]. Measurements of qE in this mutant showed that de-epoxidation of violaxanthin to zeaxanthin is necessary for most of the qE *in vivo* in leaves of *Arabidopsis*. Similar results were reported recently with violaxanthin de-epoxidase antisense tobacco plants [17]. *Arabidopsis lutein deficient 2* (*lut2*) mutants that lack lutein exhibit a partial defect in qE [18] and *npq1; lut2* double mutants completely lack qE [19], suggesting a possible role for lutein in qE. Transgenic *Arabidopsis* plants with elevated lutein have been generated by overexpression of lycopene ϵ -cyclase, resulting in an increase in

qE [20]. It is possible that lutein has a direct role in the qE mechanism or that the absence of lutein could affect qE indirectly, perhaps through a perturbation of LHC assembly and structure [21].

To identify factors besides lumen pH and xanthophylls that are necessary for qE, mutants have been isolated that lack qE but have normal pigment composition [15,16••]. In particular, the *Arabidopsis npq4* mutant is completely defective in qE and also lacks the conformational change that is monitored by measuring the absorbance change at 535 nm in leaves [16••]. Map-based cloning revealed that the *NPQ4* gene encodes photosystem II subunit S (PsbS), an integral-membrane-protein component of PS II that is a member of the LHC-protein superfamily. Like other LHC proteins, PsbS (also known as CP22 [for 22-kiloDalton chlorophyll-protein complex]) is thought to bind Chls and xanthophylls, although the stoichiometry and affinity of this binding have not been firmly established [22]. Despite the absence of PsbS protein in the *npq4* mutant, the light harvesting and photosynthetic oxygen evolution of this mutant are not impaired, suggesting that PsbS functions specifically in qE [16••]. It has been proposed that proton binding to PsbS triggers a conformational change and ¹Chl de-excitation either in the PsbS protein itself or in neighboring LHC proteins [16••].

Despite these advances in understanding the genetics of qE, the actual biophysical mechanism of ¹Chl de-excitation in qE is still unknown. An attractive proposal has been that zeaxanthin is a quencher of excess photons through a reverse transfer of excitation energy from ¹Chl to the singlet state of zeaxanthin [23], and that the xanthophyll cycle converts a light-harvesting pigment (violaxanthin) into a quencher (zeaxanthin) [24]. Until recently, the feasibility of this proposal has been unclear, because the energy levels of the lowest singlet state of the carotenoids had not been measured directly. Using the techniques of transient absorption spectroscopy and fluorescence spectroscopy, both zeaxanthin and violaxanthin have recently been found

to have low-energy singlet states that could potentially accept energy from ¹Chl [25••,26••]. Although it is possible that zeaxanthin could be acting as a quencher, the difference in energy between zeaxanthin and violaxanthin is not sufficient to explain why zeaxanthin but not violaxanthin is involved in qE [26••]. As suggested by experiments with isolated LHC proteins, it is likely that the distinct structural features of zeaxanthin and violaxanthin are important in explaining their different roles [27,28].

Long-term strategies for excess photons

In addition to coping with excess photons during short-term fluctuations in incident sunlight, many plants are confronted with environmental stresses, often on a seasonal basis, that limit their productive utilization of light energy and necessitate long-term mechanisms to deal with excess photons. Common acclimation responses to excess light include increasing the capacity for photosynthetic electron transport and CO₂ fixation, as well as decreasing the size of the light-harvesting antennae that are associated with the photosystems. This acclimation may involve the regulation of nuclear and chloroplast gene expression by redox potential [29–31], and/or the proteolytic degradation of existing LHCs [32]. In terms of safety valves, there are also mechanisms for the sustained (rather than rapidly reversible) dissipation of excess photons. Because sustained thermal dissipation results in a decrease in the quantum efficiency of photosynthesis, these mechanisms have been collectively termed photoinhibitory quenching (qI) (Figure 1). It is possible that at least part of qI may be mechanistically similar to qE [33,34], but some types of qI do not appear to depend on lumen pH [35,36].

qI has been studied extensively in overwintering evergreen plants that retain Chl despite prolonged cold-stress conditions that limit photosynthesis. These plants exhibit persistent thermal dissipation of excess photons that is reversed only slowly, even upon warming of their leaves [37]. Interestingly, zeaxanthin accumulation is associated with qI, as with qE, and recovery from qI has been correlated with the conversion of zeaxanthin back to violaxanthin in the xanthophyll cycle [38]. A possible role for protein phosphorylation in zeaxanthin epoxidation and recovery from qI has been indicated by the effects of phosphatase inhibitors on these processes [39,40].

New insights into qI have come from spectroscopic studies of overwintering evergreen (snow gum) leaves [41••]. In their winter-acclimated state, these leaves were characterized by the presence of a novel band at 710–715 nm in their Chl fluorescence emission spectra and a short-lifetime Chl-fluorescence component, indicating that they had increased thermal dissipation. Slow recovery from qI was correlated with the loss of these spectral and lifetime components and with the epoxidation of zeaxanthin to violaxanthin. These results are consistent with the idea that winter acclimation in evergreens involves significant changes in the organization and/or composition of Chl-containing

complexes. It is interesting to note that increased levels of several proteins, including PsbS, are associated with the winter acclimation of pine trees [42].

Alternative sinks for excess electrons

In linear photosynthetic electron transport, the reducing side of PS I is a key branchpoint from which electrons can be transferred for use in the major chloroplast assimilatory reactions, namely the reduction of C, N, and S. Limited availability of acceptors, especially of CO₂, can result in an excess of electrons in the photosynthetic electron-transport chain, despite the efficient dissipation of excess photons in PS II. Overreduction of electron-transport carriers may sensitize PS II to photoinhibition [3] and increase the production of other reactive oxygen species such as superoxide (O₂^{•-}) and hydrogen peroxide (H₂O₂) [1•].

Alternative electron acceptors may serve as sinks for excess electrons, thereby helping to prevent overreduction of electron carriers. Although the contribution of alternative sinks appears to be limited under steady-state conditions, they may be especially important in fluctuating light, for example during the induction phase of photosynthesis following a dark to light transfer [43]. NADPH-dependent reduction of oxaloacetate to malate has been suggested to provide such a safety valve for excess electrons, although experiments with antisense malate dehydrogenase plants have not supported a major role for this potential electron sink under steady-state conditions [44]. Thioredoxin is an electron acceptor that has important regulatory functions in chloroplasts [45], but its capacity as a sink for excess electrons is limited.

The best characterized alternative sink for electrons is photorespiratory metabolism (Figure 1). Oxygenation of ribulose-1,5-bisphosphate by Rubisco (ribulose-1,5-bisphosphate carboxylase/oxygenase) results in the formation of one molecule of 3-phosphoglycerate and one molecule of a toxic byproduct, 2-phosphoglycolate. Photorespiratory metabolism utilizes both NADPH and ATP in reactions occurring in chloroplasts, mitochondria, and peroxisomes to recover fixed C that would otherwise be lost. Thus, although it may initially appear to be a wasteful process that is a relic of Rubisco's evolutionary history, photorespiration can help to prevent light-induced damage to the photosynthetic apparatus [46].

Photoreduction of oxygen by Photosystem I: the water–water cycle

Besides the oxygenase activity of Rubisco, O₂ can be directly reduced by PS I (Figure 1). Univalent electron transfer to O₂ results in the generation of O₂^{•-}, which can subsequently be converted to H₂O₂ by superoxide dismutase. Ascorbate peroxidase activity in the chloroplast is responsible for the reduction of H₂O₂ to H₂O; the resulting monodehydroascorbate radical can be re-reduced directly by PS I. Together, these reactions are termed the water–water cycle, because electrons generated from the oxidation of water at PS II are used to reduce O₂ to water at PS I [47•]. The water–water

cycle consumes excess electrons but contributes to generation of a ΔpH without concomitant utilization of ATP.

The electron transport capacity of the water–water cycle is unclear and the extent to which it operates *in vivo* is controversial [46,48•]. Much of the uncertainty is due to difficulty in measuring flux through the water–water cycle, because there is no net O_2 exchange. Mass spectrometric measurements of gross $^{16}O_2$ evolution and $^{18}O_2$ uptake have been used to study the contribution of the water–water cycle as a safety valve in Rubisco antisense plants [49••]. A decrease in Rubisco levels would be expected to affect both the CO_2 fixation and photorespiration sinks for excess electrons, resulting in enhanced electron flow via proposed safety valves such as the water–water cycle. However, measurements of gross $^{18}O_2$ uptake by Rubisco antisense plants under steady-state non-photorespiratory conditions and at the CO_2 compensation point did not provide any indication of increased water–water cycle activity [49••]. Whether the water–water cycle plays a more significant role under other conditions, such as during photosynthetic induction, requires further investigation.

A chlororespiratory pathway involved in photoprotection

Chlororespiration is yet another type of O_2 -dependent electron transport that appears to be possible in chloroplasts (Figure 1). Nearly 20 years ago, Bennoun [50] proposed that chloroplasts contain a respiratory chain that transfers electrons from NAD(P)H to O_2 via the plastoquinone pool. Several genes encoding subunits of an NADH-plastoquinone oxidoreductase (NDH)-complex are located in the chloroplast genomes of some plants, and the corresponding enzyme activity has been found in chloroplasts [51]. Tobacco mutants lacking the NDH complex appear normal under favorable growth conditions, but they have a subtle cyclic-electron-transport defect [52,53] and increased sensitivity to photoinhibition induced by extremely strong light [54].

Until recently, there was little evidence for the existence of a chlororespiratory oxidase. However, two groups studying the *immutans* (*im*) variegation mutant of *Arabidopsis* have recently found that the *IM* gene encodes a chloroplast protein that is similar to the alternative oxidase of mitochondria [55••,56••]. Mutations in the *IM* gene act ‘plastid autonomously’ to produce white leaf sectors in which the chloroplasts accumulate phytoene [57], a colorless carotenoid precursor that cannot protect against photooxidation. Phytoene desaturation is thought to require oxidized plastoquinone as an electron acceptor [58], suggesting that the oxidation of reduced plastoquinol by the IM protein is linked to phytoene desaturation [55••,56••].

The existence of a chloroplast alternative oxidase has also been demonstrated in studies of mutants of the unicellular green alga *Chlamydomonas reinhardtii* that lack PS I. The PS-I-less mutants are unable to perform photosynthesis, but they are viable because of the ability of *Chlamydomonas* to grow heterotrophically using acetate as a source of fixed C.

Despite the lack of PS I, mass spectrometric measurements revealed that these algal mutant cells exhibit a limited but significant PS-II-dependent O_2 evolution that is matched by equivalent O_2 uptake [59,60•]. The activity persists in double mutants lacking both PS I and the cytochrome *b₆/f* complex, and in isolated chloroplasts, suggesting the involvement of a thylakoid plastoquinone oxidase that is homologous to the *Arabidopsis* IM protein [61••]. Consistent with this hypothesis, the O_2 exchange activity of *Chlamydomonas* thylakoids was inhibited by propyl gallate, which also inhibits oxidase activity in *Escherichia coli* cells expressing the *Arabidopsis* IM protein, and a *Chlamydomonas* thylakoid protein of the expected size was detected using an antibody raised against IM [61••].

Thus, chloroplasts appear to have an alternative oxidase activity that can remove electrons from the plastoquinone pool. The oxidase function may be especially critical for efficient carotenoid biosynthesis during the early stages of chloroplast development when PS-I-dependent oxidation of plastoquinol is less efficient [55••,56••]. Carotenoid biosynthesis is also necessary in fully greened chloroplasts for the repair of photodamaged PS II [62], raising the possibility that the chlororespiratory oxidase may have a significant role in providing a safety valve for excess electrons in intense light, thereby allowing carotenoid biosynthesis to proceed when the photosynthetic electron-transport chain becomes overreduced.

Conclusions

Constitutive (i.e. the triplet valve), inducible (i.e. qE), and sustained (i.e. qI) mechanisms are involved in eliminating excess photons in the Chl-containing LHCs, whereas several oxygen-dependent electron-transport processes are potential safety valves for excess electrons. Analysis of an *Arabidopsis* mutant has uncovered a role for the PsbS protein in qE [16••], raising new questions while some old questions remain. For example, future studies will address the possible involvement of proton- and pigment-binding by PsbS, and the location of PsbS relative to other PS II subunits, but the ultimate goal is still to elucidate the structural and biophysical basis of qE. At the same time, progress can be expected in understanding the mechanism of sustained qI, which is associated with a reorganization of Chl that results in fluorescence spectral and lifetime changes [41••]. It will be interesting to compare and contrast findings for qE and qI, especially because of the changes in PsbS levels that have been observed in overwintering pine needles [42]. Although the water–water cycle does not appear to be a major contributor to dissipation of excess electrons during steady-state photosynthesis [49••], its significance under rapidly fluctuating conditions warrants further investigation. A newly discovered chloroplast alternative oxidase is important for photoprotection during chloroplast development [55••,56••] and perhaps also in mature chloroplasts. Undoubtedly, the isolation of new mutants and antisense plants will allow further assessment of the physiological significance of various safety valves for photosynthesis. However, because several processes contribute to

Handwritten note: xanthophyll cycle

the dissipation of excess photons and electrons from the photosynthetic apparatus, one must consider that the phenotypes of mutants may be affected by the compensatory engagement of alternative protective mechanisms.

Acknowledgements

I thank the Searle Scholars Program/The Chicago Community Trust, the US Department of Agriculture-National Research Initiative Competitive Grants Program, and the US Department of Energy for supporting my laboratory's research on nonphotochemical quenching. I apologize to colleagues whose relevant work I was not able to cite and/or discuss extensively because of space constraints.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Niyogi KK: **Photoprotection revisited: genetic and molecular approaches.** *Annu Rev Plant Physiol Plant Mol Biol* 1999, **50**:333-359. An overview of the chloroplast processes that are involved in photoprotection of the photosynthetic apparatus. The review emphasizes the use of mutants and antisense plants.
2. Baier M, Dietz K-J: **Alkyl hydroperoxide reductases: the way out of the oxidative breakdown of lipids in chloroplasts.** *Trends Plant Sci* 1999, **4**:166-168.
3. Melis A: **Photosystem-II damage and repair cycle in chloroplasts: what modulates the rate of photodamage *in vivo*?** *Trends Plant Sci* 1999, **4**:130-135.
4. Karpinski S, Reynolds H, Karpinska B, Wingsle G, Creissen G, Mullineaux P: **Systemic signalling and acclimation in response to excess excitation energy in *Arabidopsis*.** *Science* 1999, **284**:654-657.
5. Green BR, Durnford DG: **The chlorophyll-carotenoid proteins of oxygenic photosynthesis.** *Annu Rev Plant Physiol Plant Mol Biol* 1996, **47**:685-714.
6. Foyer CH, Harbinson J: **Relationships between antioxidant metabolism and carotenoids in the regulation of photosynthesis.** In *The Photochemistry of Carotenoids*. Edited by Frank HA, Young AJ, Britton G, Cogdell RJ. Dordrecht: Kluwer Academic Publishers; 1999:305-325.
7. Hideg E, Kálai T, Hideg K, Vass I: **Photoinhibition of photosynthesis *in vivo* results in singlet oxygen production detection via nitroxide-induced fluorescence quenching in broad bean leaves.** *Biochemistry* 1998, **37**:11405-11411.
8. Horton P, Ruban AV, Walters RG: **Regulation of light harvesting in green plants.** *Annu Rev Plant Physiol Plant Mol Biol* 1996, **47**:655-684.
9. Yamamoto HY, Bugos RC, Hieber AD: **Biochemistry and molecular biology of the xanthophyll cycle.** In *The Photochemistry of Carotenoids*. Edited by Frank HA, Young AJ, Britton G, Cogdell RJ. Dordrecht: Kluwer Academic Publishers; 1999:293-303.
10. Demmig-Adams B, Adams WW III: **Xanthophyll cycle and light stress in nature: uniform response to excess direct sunlight among higher plant species.** *Planta* 1996, **198**:460-470.
11. Gilmore AM, Hazlett TL, Govindjee: **Xanthophyll cycle-dependent quenching of photosystem II chlorophyll a fluorescence: formation of a quenching complex with a short fluorescence lifetime.** *Proc Natl Acad Sci USA* 1995, **92**:2273-2277.
12. Niyogi KK, Björkman O, Grossman AR: ***Chlamydomonas* xanthophyll cycle mutants identified by video imaging of chlorophyll fluorescence quenching.** *Plant Cell* 1997, **9**:1369-1380.
13. Niyogi KK, Grossman AR, Björkman O: ***Arabidopsis* mutants define a central role for the xanthophyll cycle in the regulation of photosynthetic energy conversion.** *Plant Cell* 1998, **10**:1121-1134.
14. Shikanai T, Munekage Y, Shimizu K, Endo T, Hashimoto T: **Identification and characterization of *Arabidopsis* mutants with reduced quenching of chlorophyll fluorescence.** *Plant Cell Physiol* 1999, **40**:1134-1142.
15. Peterson RB, Havir EA: **A nonphotochemical-quenching-deficient mutant of *Arabidopsis thaliana* possessing normal pigment composition and xanthophyll-cycle activity.** *Planta* 2000, **210**:205-214.
16. Li X-P, Björkman O, Shih C, Grossman AR, Rosenquist M, Jansson S, Niyogi KK: **A pigment-binding protein essential for regulation of photosynthetic light harvesting.** *Nature* 2000, **403**:391-395. The isolation and characterization of an *Arabidopsis* mutant, *npq4-1*, which lacks qE but not the xanthophyll cycle, showed that the gene encoding PsbS is necessary for qE. Although PsbS is a member of the LHC superfamily, the protein appears to have a specialized function in the dissipation of excess photons rather than in light harvesting. PsbS is a possible evolutionary intermediate in the evolution of LHC proteins, suggesting that the photoprotective function of this protein family preceded its light-harvesting function.
17. Chang S-H, Bugos RC, Sun W-H, Yamamoto HY: **Antisense suppression of violaxanthin de-epoxidase in tobacco does not affect plant performance in controlled growth conditions.** *Photosynth Res* 2000, in press.
18. Pogson BJ, Niyogi KK, Björkman O, DellaPenna D: **Altered xanthophyll compositions adversely affect chlorophyll accumulation and nonphotochemical quenching in *Arabidopsis* mutants.** *Proc Natl Acad Sci USA* 1998, **95**:13324-13329.
19. Niyogi KK, Shih C, Chow WS, Pogson BJ, DellaPenna D, Björkman O: **Photoprotection in a zeaxanthin- and lutein-deficient double mutant of *Arabidopsis*.** *Photosynth Res* 2000, in press.
20. Pogson BJ, Rissler HM: **Genetic manipulation of carotenoid biosynthesis and photoprotection.** *Phil Trans R Soc Lond B* 2000, in press.
21. Gilmore AM: **Xanthophyll cycle-dependent nonphotochemical quenching in photosystem II: mechanistic insights gained from *Arabidopsis thaliana* L. mutants that lack violaxanthin deepoxidase and/or lutein.** *Photosynth Res* 2000, in press.
22. Funk C, Schröder WP, Napiwotzki A, Tjus SE, Renger G, Andersson B: **The PSII-S protein of higher plants: a new type of pigment-binding protein.** *Biochemistry* 1995, **34**:11133-11141.
23. Demmig-Adams B: **Carotenoids and photoprotection in plants: a role for the xanthophyll zeaxanthin.** *Biochim Biophys Acta* 1990, **1020**:1-24.
24. Frank HA, Cua A, Chynwat V, Young A, Gosztola D, Wasielewski MR: **Photophysics of the carotenoids associated with the xanthophyll cycle in photosynthesis.** *Photosynth Res* 1994, **41**:389-395.
25. Polivka T, Herek JL, Zigmantas D, Åkerlund H-E, Sundström V: **Direct observation of the (forbidden) S₁ state in carotenoids.** *Proc Natl Acad Sci USA* 1999, **96**:4914-4917. Femtosecond transient absorption spectroscopy was used to determine the energy level of the elusive S₁ state of violaxanthin and zeaxanthin. Unexpectedly, both of these xanthophylls are potential acceptors of excitation energy from ¹Chl.
26. Frank HA, Bautista JA, Josue JS, Young AJ: **Mechanism of nonphotochemical quenching in green plants: energies of the lowest singlet states of violaxanthin and zeaxanthin.** *Biochemistry* 2000, **39**:2831-2837. The finding of low-lying S₁ states of violaxanthin and zeaxanthin [25**] was corroborated using ultrasensitive fluorescence spectroscopy. The difference in energies between these xanthophylls is too small to account for the specific role of zeaxanthin in qE.
27. Ruban AV, Phillip D, Young AJ, Horton P: **Excited-state energy level does not determine the differential effect of violaxanthin and zeaxanthin on chlorophyll fluorescence quenching in the isolated light-harvesting complex of photosystem II.** *Photochem Photobiol* 1998, **68**:829-834.
28. Wentworth M, Ruban AV, Horton P: **Chlorophyll fluorescence quenching in isolated light harvesting complexes induced by zeaxanthin.** *FEBS Lett* 2000, **471**:71-74.
29. Escoubas J-M, Lomas M, LaRoche J, Falkowski PG: **Light intensity regulation of *cab* gene transcription is signaled by the redox state of the plastoquinone pool.** *Proc Natl Acad Sci USA* 1995, **92**:10237-10241.
30. Maxwell DP, Laudenbach DE, Huner NPA: **Redox regulation of light-harvesting complex II and *cab* mRNA abundance in *Dunaliella salina*.** *Plant Physiol* 1995, **109**:787-795.
31. Pfanschmidt T, Nilsson A, Allen JF: **Photosynthetic control of chloroplast gene expression.** *Nature* 1999, **397**:625-628.
32. Yang D-H, Webster J, Adam Z, Lindahl M, Andersson B: **Induction of acclimative proteolysis of the light-harvesting chlorophyll a/b protein of photosystem II in response to elevated light intensities.** *Plant Physiol* 1998, **118**:827-834.

33. Gilmore AM, Björkman O: **Temperature-sensitive coupling and uncoupling of ATPase-mediated, nonradiative energy dissipation: similarities between chloroplasts and leaves.** *Planta* 1995, **197**:646-654.
 34. Ruban AV, Horton P: **An investigation of the sustained component of nonphotochemical quenching of chlorophyll fluorescence in isolated chloroplasts and leaves of spinach.** *Plant Physiol* 1995, **108**:721-726.
 35. Verhoeven AS, Adams WW III, Demmig-Adams B: **Two forms of sustained xanthophyll cycle-dependent energy dissipation in overwintering *Euonymus kiautschovicus*.** *Plant Cell Environ* 1998, **21**:893-903.
 36. Verhoeven AS, Adams WW III, Demmig-Adams B: **The xanthophyll cycle and acclimation of *Pinus ponderosa* and *Malva neglecta* to winter stress.** *Oecologia* 1999, **118**:277-287.
 37. Adams WW III, Demmig-Adams B, Verhoeven AS, Barker DH: **'Photoinhibition' during winter stress: involvement of sustained xanthophyll cycle-dependent energy dissipation.** *Aust J Plant Physiol* 1994, **22**:261-276.
 38. Verhoeven AS, Adams WW III, Demmig-Adams B: **Close relationship between the state of the xanthophyll cycle pigments and photosystem II efficiency during recovery from winter stress.** *Physiol Plant* 1996, **96**:567-576.
 39. Xu CC, Jeon YA, Hwang HJ, Lee C-H: **Suppression of zeaxanthin epoxidation by chloroplast phosphatase inhibitors in rice leaves.** *Plant Sci* 1999, **146**:27-34.
 40. Demmig-Adams B, Adams WW III, Ebbert V, Logan BA: **Ecophysiology of the xanthophyll cycle.** In *The Photochemistry of Carotenoids*. Edited by Frank HA, Young AJ, Britton G, Cogdell RJ. Dordrecht: Kluwer Academic Publishers; 1999:245-269.
 41. Gilmore AM, Ball MC: **Protection and storage of chlorophyll in overwintering evergreens.** *Proc Natl Acad Sci USA* 2000, in press. Spectroscopic analyses showed that sustained dissipation of excess photons is associated with diagnostic changes in Chl fluorescence lifetimes, which are similar to those occurring during qE. These changes reflect the dramatic reorganization of Chl that occurs in overwintering leaves.
 42. Ottander C, Campbell D, Öquist G: **Seasonal changes in photosystem II organisation and pigment composition in *Pinus sylvestris*.** *Planta* 1995, **197**:176-183.
 43. Ruuska SA, von Caemmerer S, Badger MR, Andrews TJ, Price GD, Robinson SA: **Xanthophyll cycle, light energy dissipation and electron transport in transgenic tobacco with reduced carbon assimilation capacity.** *Aust J Plant Physiol* 2000, **27**:289-300.
 44. Backhausen JE, Emmerlich A, Holtgreffe S, Horton P, Nast G, Rogers JJM, Müller-Röber B, Scheibe R: **Transgenic potato plants with altered expression levels of chloroplast NADP-malate dehydrogenase: interactions between photosynthetic electron transport and malate metabolism in leaves and in isolated intact chloroplasts.** *Planta* 1998, **207**:105-114.
 45. Buchanan BB: **Regulation of CO₂ assimilation in oxygenic photosynthesis: the ferredoxin/thioredoxin system.** *Arch Biochem Biophys* 1991, **228**:1-9.
 46. Osmond CB, Grace SC: **Perspectives on photoinhibition and photorespiration in the field: quintessential inefficiencies of the light and dark reactions of photosynthesis?** *J Exp Botany* 1995, **46**:1351-1362.
 47. Asada K: **The water-water cycle in chloroplasts: scavenging of active oxygens and dissipation of excess photons.** *Annu Rev Plant Physiol Plant Mol Biol* 1999, **50**:601-639.
- A comprehensive summary of the biochemistry and physiological functions of the water-water cycle, which is also known as the Mehler-ascorbate-peroxidase pathway or pseudocyclic electron transport.
48. Foyer CH, Noctor G: **Oxygen processing in photosynthesis: regulation and signalling.** *New Phytol* 2000, **146**:359-388.
- A thorough and critical look at several oxygen-dependent processes that are related to photosynthesis and photoprotection.
49. Ruuska SA, Badger MR, Andrews TJ, von Caemmerer S: **Photosynthetic electron sinks in transgenic tobacco with reduced amounts of Rubisco: little evidence for significant Mehler reaction.** *J Exp Botany* 2000, **51**:357-368.
- Measurements of Chl fluorescence, CO₂ assimilation, and oxygen exchange were made in antisense tobacco plants with decreased Rubisco concentrations. Despite the expectation that alternative electron sinks would be necessary when CO₂ fixation and photorespiration are restricted, the results were not consistent with increased electron flux via the water-water cycle.
50. Bennoun P: **Evidence for a respiratory chain in the chloroplast.** *Proc Natl Acad Sci USA* 1982, **79**:4352-4356.
 51. Sazanov LA, Burrows PA, Nixon PJ: **The plastid *ndh* genes code for an NADH-specific dehydrogenase: isolation of a complex I analogue from pea thylakoid membranes.** *Proc Natl Acad Sci USA* 1998, **95**:1319-1324.
 52. Burrows PA, Sazanov LA, Svab Z, Maliga P, Nixon PJ: **Identification of a functional respiratory complex in chloroplasts through analysis of tobacco mutants containing disrupted plastid *ndh* genes.** *EMBO J* 1998, **17**:868-876.
 53. Shikanai T, Endo T, Hashimoto T, Yamada Y, Asada K, Yokota A: **Directed disruption of the tobacco *ndhB* gene impairs cyclic electron flow around photosystem I.** *Proc Natl Acad Sci USA* 1998, **95**:9705-9709.
 54. Endo T, Shikanai T, Takabayashi A, Asada K, Sato F: **The role of chloroplastic NAD(P)H dehydrogenase in photoprotection.** *FEBS Lett* 1999, **457**:5-8.
 55. Wu D, Wright DA, Wetzel C, Voytas DF, Rodermel S: **The *IMMUTANS* variegation locus of *Arabidopsis* defines a mitochondrial alternative oxidase homolog that functions during early chloroplast biogenesis.** *Plant Cell* 1999, **11**:43-55.
- Map-based cloning revealed that the *IMMUTANS* gene encodes an alternative oxidase that is involved in carotenoid biosynthesis. A model was proposed to explain the variegation phenotype of null mutants in which 'a developmental race' occurs during chloroplast biogenesis between photooxidation and inefficient phytoene desaturation.
56. Carol P, Stevenson D, Bisanz C, Breitenbach J, Sandmann G, Mache R, Coupland G, Kuntz M: **Mutations in the *Arabidopsis* gene *IMMUTANS* cause a variegated phenotype by inactivating a chloroplast terminal oxidase associated with phytoene desaturation.** *Plant Cell* 1999, **11**:57-68.
- A transposon-tagged mutant was used to clone the *IMMUTANS* gene. The IM precursor protein was shown to be imported into isolated chloroplasts, and the mature protein was associated with the thylakoids.
57. Wetzel CM, Jiang C-Z, Meehan LJ, Voytas DF, Rodermel SR: **Nuclear-organelle interactions: the *immutans* variegation mutant of *Arabidopsis* is plastid autonomous and impaired in carotenoid biosynthesis.** *Plant J* 1994, **6**:161-175.
 58. Norris SR, Barrette TR, DellaPenna D: **Genetic dissection of carotenoid synthesis in *Arabidopsis* defines plastoquinone as an essential component of phytoene desaturation.** *Plant Cell* 1995, **7**:2139-2149.
 59. Cournac L, Redding K, Bennoun P, Peltier G: **Limited photosynthetic electron flow but no CO₂ fixation in *Chlamydomonas* mutants lacking photosystem I.** *FEBS Lett* 1997, **416**:65-68.
 60. Redding K, Cournac L, Vassiliev IR, Golbeck JH, Peltier G, Rochaix J-D: **Photosystem I is indispensable for photoautotrophic growth, CO₂ fixation, and H₂ photoproduction in *Chlamydomonas reinhardtii*.** *J Biol Chem* 1999, **274**:10466-10473.
- Oxygen evolution coupled to respiration was observed in mutants that completely lack PS I due to deletions in genes encoding key reaction-center proteins. This finding and those reported in [59] disproved the idea that PS I is not necessary for photosynthetic growth and CO₂ fixation.
61. Cournac L, Redding K, Ravenel J, Rumeau D, Josse E-M, Kuntz M, Peltier G: **Electron flow between photosystem II and oxygen in chloroplasts of photosystem I-deficient algae is mediated by a quinol oxidase involved in chlororespiration.** *J Biol Chem* 2000, **275**:17256-17262.
- Light-dependent O₂ evolution and uptake observed in mutants lacking PS I was also shown to occur in isolated chloroplasts and thylakoids. *E. coli* cells expressing the *Arabidopsis* *IMMUTANS* gene had a cyanide-resistant oxidase activity that was blocked by propyl gallate. Together, the inhibition of the chlororespiratory activity of *Chlamydomonas* thylakoids by propyl gallate and the detection of a thylakoid protein that cross-reacted with an antibody raised against the IM protein suggest that the chlororespiratory O₂ uptake is due to a *Chlamydomonas* homolog of IM.
62. Trebst A, Depka B: **Role of carotene in the rapid turnover and assembly of photosystem II in *Chlamydomonas reinhardtii*.** *FEBS Lett* 1997, **400**:359-362.